



Self-Regulation of Blood Oxygenation Level Dependent Response: Primary Effect or Epiphenomenon?

Andrea Caria^{1,2*}

¹ Department of Psychology and Cognitive Science, University of Trento, Rovereto, Italy, ² Institute of Medical Psychology and Behavioral Neurobiology, Eberhard-Karls-University of Tübingen, Tübingen, Germany

Keywords: real-time fMRI, BOLD signal, neurofeedback, self-regulation

OPEN ACCESS

Edited by:

Manuel Fernando Casanova,
University of Louisville, USA

Reviewed by:

Hari S. Sharma,
Uppsala University, Sweden
Victor Manuel Pulgar,
Wake Forest School of Medicine, USA

*Correspondence:

Andrea Caria
andrea.caria@unitn.it

Specialty section:

This article was submitted to
Neural Technology,
a section of the journal
Frontiers in Neuroscience

Received: 28 January 2016

Accepted: 09 March 2016

Published: 24 March 2016

Citation:

Caria A (2016) Self-Regulation of
Blood Oxygenation Level Dependent
Response: Primary Effect or
Epiphenomenon?
Front. Neurosci. 10:117.
doi: 10.3389/fnins.2016.00117

In the last decade, several studies indicated that neuronal activity can be volitionally modulated using real-time fMRI (rtfMRI) based neurofeedback. Human participants through rtfMRI paradigms can learn to regulate the blood oxygenation level dependent (BOLD) response in several localized cortical and subcortical regions (for extended reviews see Caria et al., 2012; Weiskopf, 2012; Sulzer et al., 2013). Increasing evidence also indicated that strengthening or weakening specific BOLD activity using rtfMRI training leads to significant changes in cognitive, emotional and motor behavior (Caria et al., 2012; Weiskopf, 2012; Sulzer et al., 2013). These findings suggested that rtfMRI might represent an important novel approach in cognitive neuroscience by potentially providing indications of cause-and-effect relationships between brain and behavior, and also in clinical applications (Subramanian et al., 2011; Linden et al., 2012; Ruiz et al., 2013; Sitaram et al., 2014).

Although rtfMRI studies extend and enrich a large body of literature demonstrating operant learning of neuronal oscillations, skepticism still exists on the validity and reliability of studies showing learned control of the BOLD response. In particular, it is still debated whether this phenomenon is a primary effect of learning or it just an epiphenomenon resulting, for instance, from repeated execution of some mental processes.

Here, I will discuss specific psychophysiological and neurophysiological mechanisms presumably underlying learned regulation of the BOLD response to attempt to clarify its nature.

OPERANT LEARNING OF THE BOLD RESPONSE

Direct manipulation of brain activity grounds on the principles of operant conditioning that describe the relationship between changes in behavior (in this case the neurophysiological response) as consequence of contingent environmental events. Reinforcement is the mechanism that allows participants to increase the frequency of specific brain activity and to shape the desired pattern of neuronal activations. During operant learning of the BOLD response participants are reinforced proportionally to how much the ongoing metabolic signal approach or resemble the target level of activation in single or multiple regions of interest. The information of the ongoing signal fluctuations, provided as explicit visual feedback to the participants, represents the intrinsic reward.

Operant learning is an important approach for manipulating brain functions (Birbaumer and Kimmel, 1979) acting on intrinsic physiological properties of neuronal activity, which have been extensively shown to undergo classical and operant conditioning (Kandel and Schwartz, 1982).

The application of operant control principles to metabolic signals builds on several previous experiments demonstrating operant control of neuroelectric activity in non-human and human

animals (Birbaumer and Kimmel, 1979; Fetz, 2007). Studies on volitional control of neural activity dates back to the late sixties and seventies when operant conditioning of central nervous system activity was demonstrated (Olds, 1965; Fetz, 1969; Shinkman et al., 1974). CNS unit conditioning was shown by operantly rewarding rats to increase the activity of midbrain neurons using intracranial stimulation (Olds, 1965). In 1969, Fetz demonstrated conditioning of the activity of single neurons in precentral cortex in anesthetized monkeys by reinforcing high rates of neuronal discharge with food reward and auditory or visual feedback of unit firing rates (Fetz, 1969). More recently, Brain-Computer Interface and Brain-Machine interface studies reported volitional control of cortical cell activity, in particular in the motor cortex, in both animals and humans (Serruya et al., 2002; Taylor et al., 2002; Carmena et al., 2003; Moritz et al., 2008), with food reward and visual feedback of neuronal ensembles' activity. Altogether these findings robustly proved that neuronal activity from localized brain areas can be manipulated through operant conditioning.

On the other hand, there are unclear aspects of operant conditioning of the BOLD response in humans that make rtfMRI neurofeedback paradigms not fully established. Among these is the role of mental and cognitive processes for learning. Some rtfMRI studies indicated that successful BOLD regulation relies on the use of mental imagery (Chiew et al., 2012; Banca et al., 2015; Blefari et al., 2015). In particular, these studies suggested that strategies based on mental imagery are important for learning BOLD control considering the partial common substrates for internal representations and overt behavior, which holds in particular for the primary motor (Jeannerod, 1995; Roth et al., 1996; Jeannerod and Frak, 1999; Niyazov et al., 2005) and visual areas (Kosslyn and Thompson, 2003; Klein et al., 2004; Slotnick et al., 2005). On the contrary, other studies indicated that a combination of cognitive strategies (mental imagery) and feedback information is critical for participants to learn BOLD regulation, and that cognitive and imagery alone is not sufficient (decharms et al., 2005; Bray et al., 2007; Caria et al., 2010; Scharnowski et al., 2012). In particular, mere repetition of mental strategies leads to an initial increase of BOLD signal but to a rapid decrease in the following consecutive runs (Caria et al., 2010). In fact, fMRI adaptation paradigms (Kourtzi and Kanwisher, 2001) show that task repetition leads to a decrease of the BOLD signal.

In addition, while it is true that imagery and cognitive processing certainly affect learning and physiological regulation, studies on operant control of brain oscillations suggest that feedback is more important than instructions for successful slow cortical potentials regulation (Roberts et al., 1989; Birbaumer et al., 1990). There are indications from EEG based neurofeedback studies that mental imagery is efficacious initially but it would be then dropped when participants become more confident with self-regulation (Leuthardt et al., 2004). Ultimately, control of neuroelectric activity turns to a highly implicit process over training, and the contribution of higher order cognitive processes becomes negligible. Indeed, control of neurophysiological signals can even be attained when participants have little or no direct experience of it (Rockstroh et al., 1984). However, until now there exist no clear

demonstrations that these mechanisms, although plausible, hold also for learned metabolic signals.

In most rtfMRI studies participants were aware of the responses leading to reward. Only one study reported to have employed implicit visual task and feedback to learn BOLD control in V1/V2 areas, however participants still reported to have used explicit mental strategies (Shibata et al., 2011).

In short, the fact that operant control of brain activity, even single cell responses is possible also in animals speaks against a fundamental influence of cognitive factors on the effect of operant brain regulation. On the other hand, it is conceivable that in human participants the combination of specific conscious processes and operant strategies might support retention of learned BOLD control out of the laboratory setting, and might facilitate the (re-)activation of impaired or dormant mechanisms, in particular in patients.

WHAT NEUROPHYSIOLOGICAL MECHANISMS MEDIATE LEARNED BOLD CONTROL?

A further issue of rtfMRI experiments is the unclear neurophysiology of the BOLD signal, which prevents us to univocally interpret the behavioral changes induced by operant learning of the metabolic signal. We know from previous investigations on instrumental learning of neuroelectric signals that the brain has intrinsic neurophysiological processes allowing regulation of neural activity even in the absence of external stimuli (Wolpaw et al., 2002; Birbaumer and Cohen, 2007; Fetz, 2007). While no specific receptors directly support regulation of brain oscillations, it exists a complex neurovascular system that regulates and controls the BOLD signal.

At least three main blood flow regulatory mechanisms have been described: cerebral autoregulation—the brain ability to maintain a constant flow through changes of cerebral perfusion pressure—neurogenic regulation—the cerebral blood flow modulation through extensive arborization of perivascular nerves—and flow-metabolism coupling, or functional hyperemia—the brain capacity to vary blood flow to match metabolic activity (Peterson et al., 2011). Endothelial cells and astrocytes play a central role in all these regulatory mechanisms (Iadecola and Nedergaard, 2007): the former by providing several vasoactive factors for the regulation of cerebral blood flow and the latter because of their anatomical position that physically links the cerebral microvasculature with synapses.

Learned increase and decrease of BOLD activity might indeed be associated with variations in the flow-metabolism coupling. Local changes in neural activity have been shown to influence BOLD signal (Buxton et al., 2004; Logothetis and Wandell, 2004; Lee et al., 2010) through variations in cerebral blood flow (CBF), cerebral blood volume (CBV) and cerebral metabolic rate of oxygen consumption (CMRO₂). It has been also demonstrated that changes in the coupling ratio of CBF and CMRO₂ in response to neural activity strongly affect the BOLD response (Kim and Ogawa, 2012; Buxton et al., 2014). Considering such complexity of the neurovascular system, it is thus conceivable

that learned control of the BOLD response might be based on these sophisticated regulatory mechanisms.

An intriguing perspective for rtfMRI neurofeedback arises from the hypothesis proposing that modulation of brain activity is directly and causally affected by changes of metabolic signals (Moore and Cao, 2008). Specifically, it has been conjectured that the brain vascular system, being a complex and interconnected network under tight regulatory control that occurs in close communication with neurons and glia, might directly contribute to information processing (Moore and Cao, 2008). Accordingly, the hemodynamic processes would not only support metabolic demand but also directly shape brain functions; self-regulation of metabolic signals might thus induce changes of neuronal activations. In line with this perspective non-invasive direct electrical stimulation (e.g., using transcranial direct electric stimulation, tDCS) targeting cerebral microvessels has been proposed to enhance brain functions through changes of cerebrovascular function (Dutta, 2015; Pulgar, 2015).

So far, despite the increasing number of hypotheses and studies aiming to clarify the nature of the BOLD response, the neurometabolic coupling of this signal remains unclear. It has been shown that the BOLD fMRI response correlates with local field potentials (Logothetis et al., 2001; Viswanathan and Freeman, 2007), with spiking activity (Heeger and Ress, 2002), and with both LFP and spiking activity (Mukamel et al., 2005). In addition, BOLD signal can reflect different brain processing such as excitatory and inhibitory activity, neuromodulation, and bottom-up and top-down signals (Viswanathan and Freeman, 2007; Logothetis, 2008; Lee et al., 2010).

On the basis of this evidence it is difficult to infer whether the net effect of self-regulation of BOLD signal is excitatory, inhibitory or a combination of both. On the other hand, rtfMRI studies, although sometimes controversial, often reported an improvement of subjects' performance associated with increased amplitude of the BOLD response (decharms et al., 2005; Bray et al., 2007; Rota et al., 2009; Caria et al., 2010; Scharnowski et al., 2012, 2015; Zhang et al., 2013; Blefari et al., 2015). For instance, studies showed a decrease in reaction times for task execution associated with increase activity of the primary motor cortex (M1), and for task initiation after up-regulation of the SMA (Scharnowski et al., 2015). M1 activity was also positively correlated with accuracy improvement of performance during an isometric pinching task (Blefari et al., 2015). In the visual domain, increased visual perception was induced by increased BOLD response in the visual cortex (Scharnowski et al., 2012). Moreover, enhanced emotional response was associated with increased percentage BOLD signal change in the left anterior insula (Caria et al., 2010). Importantly, these studies reported poorer performance either during initial regulation runs or after down-regulation. Altogether these findings suggest that learned increase of the BOLD response might be associated with increased excitatory activity. This assumption would be also supported by studies showing improved memory performance

associated with decreased BOLD activity in the parahippocampal cortex (Yoo et al., 2012; Scharnowski et al., 2015), which was then interpreted as result of increased brain resources for memory encoding because of decreased cortical processing (Yoo et al., 2012). Indeed, increased BOLD signal might occur from increased net excitation associated with glutamatergic activity (Logothetis, 2008). However, this represents only one possible interpretation, as larger BOLD signal can also be induced by concurrent increase of both glutamatergic and GABAergic activity, with changes in hemodynamic responses reflecting balanced local recurrent activity (Logothetis, 2008). In line with this alternative mechanism, it has been shown that enhanced perceptual sensitivity induced by fMRI feedback training results from a mixture of positive and negative activated voxels in the targeted regions of interest rather than from uniform positive or negative activity (Shibata et al., 2011).

Therefore, fMRI neurofeedback might also rely on, and possibly alter, the capacity of the cerebral cortex to generate persistent activity in the absence of sensory stimulation (Haider et al., 2006), which would in turn influence neuronal responsiveness to a wide range of inputs (Shu et al., 2003a). In addition, as the ongoing network activity can be turned on and off by synaptic inputs via electrical stimulation (Shu et al., 2003b), it can also be postulated its self-regulation via operant conditioning.

CONCLUSIONS

In short, the observed behavioral changes induced by rtfMRI training might be related to several plausible neurophysiological processes, and it is still not possible to conclusively ascertain the actual mechanism on the basis of the current results. Multimodal measurements such as simultaneous EEG-rtfMRI acquisition, as well as alternative real-time functional imaging methods such as MR perfusion and MR spectroscopy, might help to clarify the neurophysiological mechanisms underlying learned control of the BOLD response. Nevertheless, there exist clear indications that self-regulation of BOLD signal is not an epiphenomenon but a primary effect of operant learning, which can rely on precise neurovascular regulatory mechanisms.

AUTHOR CONTRIBUTIONS

The author confirms being the sole contributor of this work and approved it for publication.

ACKNOWLEDGMENTS

This work is part of the BRAINTRAIN European research network (Collaborative Project) supported by the European Commission under the Health Cooperation Work Programme of the 7th Framework Programme (Grant agreement n° 602186).

REFERENCES

- Banca, P., Sousa, T., Catarina Duarte, I., and Castelo-Branco, M. (2015). Visual motion imagery neurofeedback based on the hMT+/V5 complex: evidence for a feedback-specific neural circuit involving neocortical and cerebellar regions. *J. Neural Eng.* 12:066003. doi: 10.1088/1741-2560/12/6/066003
- Birbaumer, N., and Cohen, L. G. (2007). Brain-computer interfaces: communication and restoration of movement in paralysis. *J. Physiol.* 579, 621–636. doi: 10.1113/jphysiol.2006.125633
- Birbaumer, N., Elbert, T., Canavan, A. G., and Rockstroh, B. (1990). Slow potentials of the cerebral cortex and behavior. *Physiol. Rev.* 70, 1–41.
- Birbaumer, N., and Kimmel, H. (1979). *Biofeedback and Self-Regulation*. Hillsdale, NJ: Erlbaum.
- Blefari, M. L., Sulzer, J., Hepp-Reymond, M. C., Kollias, S., and Gassert, R. (2015). Improvement in precision grip force control with self-modulation of primary motor cortex during motor imagery. *Front. Behav. Neurosci.* 9:18. doi: 10.3389/fnbeh.2015.00018
- Bray, S., Shimojo, S., and O'doherty, J. P. (2007). Direct instrumental conditioning of neural activity using functional magnetic resonance imaging-derived reward feedback. *J. Neurosci.* 27, 7498–7507. doi: 10.1523/JNEUROSCI.2118-07.2007
- Buxton, R. B., Griffeth, V. E., Simon, A. B., Moradi, F., and Shmuel, A. (2014). Variability of the coupling of blood flow and oxygen metabolism responses in the brain: a problem for interpreting BOLD studies but potentially a new window on the underlying neural activity. *Front. Neurosci.* 8:139. doi: 10.3389/fnins.2014.00139
- Buxton, R. B., Uludag, K., Dubowitz, D. J., and Liu, T. T. (2004). Modeling the hemodynamic response to brain activation. *Neuroimage* 23(Suppl. 1), S220–S233. doi: 10.1016/j.neuroimage.2004.07.013
- Caria, A., Sitaram, R., and Birbaumer, N. (2012). Real-time fMRI: a tool for local brain regulation. *Neuroscientist* 18, 487–501. doi: 10.1177/1073858411407205
- Caria, A., Sitaram, R., Veit, R., Begliomini, C., and Birbaumer, N. (2010). Volitional control of anterior insula activity modulates the response to aversive stimuli. A real-time functional magnetic resonance imaging study. *Biol. Psychiatry* 68, 425–432. doi: 10.1016/j.biopsych.2010.04.020
- Carmena, J. M., Lebedev, M. A., Crist, R. E., O'doherty, J. E., Santucci, D. M., Dimitrov, D. F., et al. (2003). Learning to control a brain-machine interface for reaching and grasping by primates. *PLoS Biol.* 1:E42. doi: 10.1371/journal.pbio.0000042
- Chiew, M., Laconte, S. M., and Graham, S. J. (2012). Investigation of fMRI neurofeedback of differential primary motor cortex activity using kinesthetic motor imagery. *Neuroimage* 61, 21–31. doi: 10.1016/j.neuroimage.2012.02.053
- decharms, R. C., Maeda, F., Glover, G. H., Ludlow, D., Pauly, J. M., Soneji, D., et al. (2005). Control over brain activation and pain learned by using real-time functional MRI. *Proc. Natl. Acad. Sci. U.S.A.* 102, 18626–18631. doi: 10.1073/pnas.0505210102
- Dutta, A. (2015). Bidirectional interactions between neuronal and hemodynamic responses to transcranial direct current stimulation (tDCS): challenges for brain-state dependent tDCS. *Front. Syst. Neurosci.* 9:107. doi: 10.3389/fnsys.2015.00107
- Fetz, E. E. (1969). Operant conditioning of cortical unit activity. *Science* 163, 955–958. doi: 10.1126/science.163.3870.955
- Fetz, E. E. (2007). Volitional control of neural activity: implications for brain-computer interfaces. *J. Physiol.* 579, 571–579. doi: 10.1113/jphysiol.2006.127142
- Haider, B., Duque, A., Hasenstaub, A. R., and McCormick, D. A. (2006). Neocortical network activity *in vivo* is generated through a dynamic balance of excitation and inhibition. *J. Neurosci.* 26, 4535–4545. doi: 10.1523/JNEUROSCI.5297-05.2006
- Heeger, D. J., and Ress, D. (2002). What does fMRI tell us about neuronal activity? *Nat. Rev. Neurosci.* 3, 142–151. doi: 10.1038/nrn730
- Iadecola, C., and Nedergaard, M. (2007). Glial regulation of the cerebral microvasculature. *Nat. Neurosci.* 10, 1369–1376. doi: 10.1038/nn2003
- Jeannerod, M. (1995). Mental imagery in the motor context. *Neuropsychologia* 33, 1419–1432. doi: 10.1016/0028-3932(95)00073-C
- Jeannerod, M., and Frak, V. (1999). Mental imaging of motor activity in humans. *Curr. Opin. Neurobiol.* 9, 735–739. doi: 10.1016/S0959-4388(99)00038-0
- Kandel, E. R., and Schwartz, J. H. (1982). Molecular biology of learning: modulation of transmitter release. *Science* 218, 433–443. doi: 10.1126/science.6289442
- Kim, S. G., and Ogawa, S. (2012). Biophysical and physiological origins of blood oxygenation level-dependent fMRI signals. *J. Cereb. Blood Flow Metab.* 32, 1188–1206. doi: 10.1038/jcbfm.2012.23
- Klein, I., Dubois, J., Mangin, J. F., Kherif, F., Flandin, G., Poline, J. B., et al. (2004). Retinotopic organization of visual mental images as revealed by functional magnetic resonance imaging. *Brain Res. Cogn. Brain Res.* 22, 26–31. doi: 10.1016/j.cogbrainres.2004.07.006
- Kosslyn, S. M., and Thompson, W. L. (2003). When is early visual cortex activated during visual mental imagery? *Psychol. Bull.* 129, 723–746. doi: 10.1037/0033-2909.129.5.723
- Kourtzi, Z., and Kanwisher, N. (2001). Representation of perceived object shape by the human lateral occipital complex. *Science* 293, 1506–1509. doi: 10.1126/science.1061133
- Lee, J. H., Durand, R., Gradinaru, V., Zhang, F., Goshen, I., Kim, D. S., et al. (2010). Global and local fMRI signals driven by neurons defined optogenetically by type and wiring. *Nature* 465, 788–792. doi: 10.1038/nature09108
- Leuthardt, E. C., Schalk, G., Wolpaw, J. R., Ojemann, J. G., and Moran, D. W. (2004). A brain-computer interface using electrocorticographic signals in humans. *J. Neural Eng.* 1, 63–71. doi: 10.1088/1741-2560/1/2/001
- Linden, D. E., Habes, I., Johnston, S. J., Linden, S., Tatineni, R., Subramanian, L., et al. (2012). Real-time self-regulation of emotion networks in patients with depression. *PLoS ONE* 7:e38115. doi: 10.1371/journal.pone.0038115
- Logothetis, N. K. (2008). What we can do and what we cannot do with fMRI. *Nature* 453, 869–878. doi: 10.1038/nature06976
- Logothetis, N. K., Pauls, J., Augath, M., Trinath, T., and Oeltermann, A. (2001). Neurophysiological investigation of the basis of the fMRI signal. *Nature* 412, 150–157. doi: 10.1038/35084005
- Logothetis, N. K., and Wandell, B. A. (2004). Interpreting the BOLD signal. *Annu. Rev. Physiol.* 66, 735–769. doi: 10.1146/annurev.physiol.66.082602.092845
- Moore, C. I., and Cao, R. (2008). The hemo-neural hypothesis: on the role of blood flow in information processing. *J. Neurophysiol.* 99, 2035–2047. doi: 10.1152/jn.01366.2006
- Moritz, C. T., Perlmuter, S. I., and Fetz, E. E. (2008). Direct control of paralysed muscles by cortical neurons. *Nature* 456, 639–642. doi: 10.1038/nature07418
- Mukamel, R., Gelbard, H., Arieli, A., Hasson, U., Fried, I., and Malach, R. (2005). Coupling between neuronal firing, field potentials, and FMRI in human auditory cortex. *Science* 309, 951–954. doi: 10.1126/science.1110913
- Niyazov, D. M., Butler, A. J., Kadah, Y. M., Epstein, C. M., and Hu, X. P. (2005). Functional magnetic resonance imaging and transcranial magnetic stimulation: effects of motor imagery, movement and coil orientation. *Clin. Neurophysiol.* 116, 1601–1610. doi: 10.1016/j.clinph.2005.02.028
- Olds, J. (1965). Operant conditioning of single unit responses. *Excerpta Med. Int. Cong. Series* 87, 372–380.
- Peterson, E. C., Wang, Z., and Britz, G. (2011). Regulation of cerebral blood flow. *Int. J. Vasc. Med.* 2011:823525. doi: 10.1155/2011/823525
- Pulgar, V. M. (2015). Direct electric stimulation to increase cerebrovascular function. *Front. Syst. Neurosci.* 9:54. doi: 10.3389/fnsys.2015.00054
- Roberts, L. E., Birbaumer, N., Rockstroh, B., Lutzenberger, W., and Elbert, T. (1989). Self-report during feedback regulation of slow cortical potentials. *Psychophysiology* 26, 392–403. doi: 10.1111/j.1469-8986.1989.tb01941.x
- Rockstroh, B., Birbaumer, N., Elbert, T., and Lutzenberger, W. (1984). Operant control of EEG and event-related and slow brain potentials. *Biofeedback Self Regul.* 9, 139–160. doi: 10.1007/BF00998830
- Rota, G., Sitaram, R., Veit, R., Erb, M., Weiskopf, N., Dogil, G., et al. (2009). Self-regulation of regional cortical activity using real-time fMRI: the right inferior frontal gyrus and linguistic processing. *Hum. Brain Mapp.* 30, 1605–1614. doi: 10.1002/hbm.20621
- Roth, M., Decety, J., Raybaudi, M., Massarelli, R., Delon-Martin, C., Segebarth, C., et al. (1996). Possible involvement of primary motor cortex in mentally simulated movement: a functional magnetic resonance imaging study. *Neuroreport* 7, 1280–1284. doi: 10.1097/00001756-199605170-00012
- Ruiz, S., Lee, S., Soekadar, S. R., Caria, A., Veit, R., Kircher, T., et al. (2013). Acquired self-control of insula cortex modulates emotion recognition and brain network connectivity in schizophrenia. *Hum. Brain Mapp.* 34, 200–212. doi: 10.1002/hbm.21427

- Scharnowski, F., Hutton, C., Josephs, O., Weiskopf, N., and Rees, G. (2012). Improving visual perception through neurofeedback. *J. Neurosci.* 32, 17830–17841. doi: 10.1523/JNEUROSCI.6334-11.2012
- Scharnowski, F., Veit, R., Zopf, R., Studer, P., Bock, S., Diedrichsen, J., et al. (2015). Manipulating motor performance and memory through real-time fMRI neurofeedback. *Biol. Psychol.* 108, 85–97. doi: 10.1016/j.biopsycho.2015.03.009
- Serruya, M. D., Hatsopoulos, N. G., Paninski, L., Fellows, M. R., and Donoghue, J. P. (2002). Instant neural control of a movement signal. *Nature* 416, 141–142. doi: 10.1038/416141a
- Shibata, K., Watanabe, T., Sasaki, Y., and Kawato, M. (2011). Perceptual learning incepted by decoded fMRI neurofeedback without stimulus presentation. *Science* 334, 1413–1415. doi: 10.1126/science.1212003
- Shinkman, P. G., Bruce, C. J., and Pfingst, B. E. (1974). Operant conditioning of single-unit response patterns in visual cortex. *Science* 184, 1194–1196. doi: 10.1126/science.184.4142.1194
- Shu, Y., Hasenstaub, A., Badoual, M., Bal, T., and McCormick, D. A. (2003a). Barrages of synaptic activity control the gain and sensitivity of cortical neurons. *J. Neurosci.* 23, 10388–10401.
- Shu, Y., Hasenstaub, A., and McCormick, D. A. (2003b). Turning on and off recurrent balanced cortical activity. *Nature* 423, 288–293. doi: 10.1038/nature01616
- Sitaram, R., Caria, A., Veit, R., Gaber, T., Ruiz, S., and Birbaumer, N. (2014). Volitional control of the anterior insula in criminal psychopaths using real-time fMRI neurofeedback: a pilot study. *Front. Behav. Neurosci.* 8:344. doi: 10.3389/fnbeh.2014.00344
- Slotnick, S. D., Thompson, W. L., and Kosslyn, S. M. (2005). Visual mental imagery induces retinotopically organized activation of early visual areas. *Cereb. Cortex* 15, 1570–1583. doi: 10.1093/cercor/bhi035
- Subramanian, L., Hindle, J. V., Johnston, S., Roberts, M. V., Husain, M., Goebel, R., et al. (2011). Real-time functional magnetic resonance imaging neurofeedback for treatment of Parkinson's disease. *J. Neurosci.* 31, 16309–16317. doi: 10.1523/JNEUROSCI.3498-11.2011
- Sulzer, J., Haller, S., Scharnowski, F., Weiskopf, N., Birbaumer, N., Blefari, M. L., et al. (2013). Real-time fMRI neurofeedback: progress and challenges. *Neuroimage* 76, 386–399. doi: 10.1016/j.neuroimage.2013.03.033
- Taylor, D. M., Tillery, S. I., and Schwartz, A. B. (2002). Direct cortical control of 3D neuroprosthetic devices. *Science* 296, 1829–1832. doi: 10.1126/science.1070291
- Viswanathan, A., and Freeman, R. D. (2007). Neurometabolic coupling in cerebral cortex reflects synaptic more than spiking activity. *Nat. Neurosci.* 10, 1308–1312. doi: 10.1038/nn1977
- Weiskopf, N. (2012). Real-time fMRI and its application to neurofeedback. *Neuroimage* 62, 682–692. doi: 10.1016/j.neuroimage.2011.10.009
- Wolpaw, J. R., Birbaumer, N., Mcfarland, D. J., Pfurtscheller, G., and Vaughan, T. M. (2002). Brain-computer interfaces for communication and control. *Clin. Neurophysiol.* 113, 767–791. doi: 10.1016/S1388-2457(02)00057-3
- Yoo, J. J., Hinds, O., Ofen, N., Thompson, T. W., Whitfield-Gabrieli, S., Triantafyllou, C., et al. (2012). When the brain is prepared to learn: enhancing human learning using real-time fMRI. *Neuroimage* 59, 846–852. doi: 10.1016/j.neuroimage.2011.07.063
- Zhang, G., Yao, L., Zhang, H., Long, Z., and Zhao, X. (2013). Improved working memory performance through self-regulation of dorsal lateral prefrontal cortex activation using real-time fMRI. *PLoS ONE* 8:e73735. doi: 10.1371/journal.pone.0073735

Conflict of Interest Statement: The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Caria. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.